



Live Well, Work Well

Health and wellness tips for your work, home and life—brought to you by the insurance professionals at The Reschini Group

For those prone to serious allergic reactions, your doctor may prescribe an auto-injector containing epinephrine.

CHILDREN'S HEALTH: ALLERGIES

An allergy exists when one's immune system responds abnormally to a usually harmless substance. For most affected people, allergies first appear during childhood.

Risk Factors

Though any child may develop allergies, there are factors that increase the risk. Family history plays a large role in a child's tendency to have an allergy. In addition, hormones, stress, smoke, and other environmental factors can play a role.

Common Allergies

Common child allergies include:

- Allergic rhinitis (commonly known as hay fever)
 - Caused by inhaled pollen or other allergens; can be seasonal or year-round
- Atopic dermatitis (eczema) - Itchy skin rash
- Food allergies - Common ones in children are peanuts, eggs, and milk. Children sometimes outgrow these allergies.
- Medications
- Insect stings or bites

Allergic Reactions

Symptoms of an allergic reaction vary by person. For inhaled or skin allergens, common symptoms include:

- Itchy, watery eyes
- Runny nose or sneezing
- Feeling tired or ill
- Rashes or hives

Food or medication allergies can cause cramps, vomiting, swelling in the mouth or throat, itchy skin, wheezing, and hives. Insect stings often cause swelling, itching, redness, and pain.

The previous examples are the most common allergies and symptoms, but remember that any number of substances can cause allergies, with a large variety of potential symptoms.

Anaphylaxis

The most severe reaction to any allergen is called anaphylaxis. This can occur for any allergy, but children with peanut or insect sting allergies may be particularly vulnerable. Anaphylaxis is a life threatening reaction, with symptoms such as full body hives, difficulty breathing, tingling of the skin, and unconsciousness. This type of reaction occurs almost immediately after exposure to the allergen, and requires emergency medical care.

Diagnosis and Treatment

Watch for abnormal symptoms in your child, particularly if they occur after eating or taking medication, after being stung, or when exposed to potential allergens indoors or outdoors. If you suspect your child has an allergy, see a doctor. A doctor can perform skin, blood, or other tests to identify the potential allergy.

If your child is diagnosed with an allergy, learn as much as you can about it so that you can adapt your family's lifestyle to avoid allergy triggers. If your child is prone to severe reactions, alert family, friends, and your child's school or child care provider, and have your child wear a medical alert bracelet at all times.

